



References

Implant Material

1. Klinge U, Klosterhalfen B, Ottinger AP, et al (2002)
PVDF as a new polymer for the construction of surgical meshes.
Biomaterials 23:3487-3493
2. Klink CD, Junge K, Binnebösel M, et al (2011)
Comparison of long-term biocompatibility of PVDF and PP meshes.
J Invest Surg 24:292-299. <https://doi.org/10.3109/08941939.2011.589883>
3. Gerullis H, Georgas E, Eimer C, et al (2011)
Evaluation of Biocompatibility of Alloplastic Materials: Development of a Tissue Culture In Vitro Test System.
Surgical technology international 21:21
4. Gerullis H, Klosterhalfen B, Borós M, et al (2013)
IDEAL in Meshes for Prolapse, Urinary Incontinence, and Hernia Repair.
Surg Innov. <https://doi.org/10.1177/1553350612472987>
5. Laroche G, Marois Y, Schwarz E, et al (1995)
Polyvinylidene fluoride monofilament sutures: can they be used safely for long-term anastomoses in the thoracic aorta?
Artif Organs 19:1190-1199
10. Berger D, Bientzle M (2008)
Polyvinylidene fluoride: a suitable mesh material for laparoscopic incisional and parastomal hernia repair!
Hernia 13:167-172. <https://doi.org/10.1007/s10029-008-0435-4>
11. Junge K, Binnebösel M, Rosch R, et al (2008)
Adhesion formation of a polyvinylidenefluoride/polypropylene mesh for intra-abdominal placement in a rodent animal model
Surgical Endoscopy 23:327-333. <https://doi.org/10.1007/s00464-008-9923-y>
20. Göretzlehner U, Müllen A (2007)
PVDF als Implantat-Werkstoff in der Urogynäkologie.
BIOmaterialien 8 (S1):28-29
27. Mary C, Marois Y, King MW, et al (1998)
Comparison of the in vivo behavior of polyvinylidene fluoride and polypropylene sutures used in vascular surgery.
ASAIO J 44:199-206
50. Roman S, Urbánková I, Callewaert G, et al (2016)
Evaluating Alternative Materials for the Treatment of Stress Urinary Incontinence and Pelvic Organ Prolapse: A Comparison of the In Vivo Response to Meshes Implanted in Rabbits.
The Journal of Urology 196:261-269. <https://doi.org/10.1016/j.juro.2016.02.067>
52. Silva RA, Silva PA, Carvalho ME (2007)
Degradation studies of some polymeric biomaterials: Polypropylene (PP) and polyvinylidene difluoride (PVDF).
THERMEC 2006, Pts 1-5 539-543:573-576
68. Conze J, Junge K, Weiss C, et al (2008)
New polymer for intra-abdominal meshes--PVDF copolymer.
J Biomed Mater Res Part B Appl Biomater 87:321-328. <https://doi.org/10.1002/jbm.b.31106>

References

Implant Material

91. Hara T (2004)
Ten-Year Results of Anterior Chamber Fixation of the Posterior ChamberIntraocular Lens.
Arch Ophthalmol 122:1112. <https://doi.org/10.1001/archophth.122.8.1112>
93. Wang H, Klosterhalfen B, Müllen A, et al (2021)
Degradation resistance of PVDF mesh in vivo in comparison to PP mesh.
J Mech Behav Biomed Mater 119:104490. <https://doi.org/10.1016/j.jmbbm.2021.104490>
100. Karabulut A, Simavli SA, Abban GM, et al (2016)
Tissue reaction to urogynecologic meshes: effect of steroid soaking in two different mesh models.
Int Urogynecol J 27:1583–1589. <https://doi.org/10.1007/s00192-016-3013-9>

References

Product Design

6. Mühl T, Binnebösel M, Klinge U, Goedderz T (2008)
New objective measurement to characterize the porosity of textile implants.
Journal of Biomedical Materials Research Part B: Applied Biomaterials 84B:176–183. <https://doi.org/10.1002/jbm.b.30859>
8. Klinge U, Klosterhalfen B (2012)
Modified classification of surgical meshes for hernia repair based on the analyses of 1,000 explanted meshes.
Hernia 16:251–258. <https://doi.org/10.1007/s10029-012-0913-6>
25. Klosterhalfen B, Junge K, Klinge U (2005)
The lightweight and large porous mesh concept for hernia repair.
Expert Rev Med Devices 2:103–117. <https://doi.org/10.1586/17434440.2.1.103>
26. Otto J, Kaldenhoff E, Kirschner-Hermanns R, et al (2013)
Elongation of textile pelvic floor implants under load is related to complete loss of effective porosity, thereby favouring incorporation in scar plates.
Journal of Biomedical Materials Research Part A n/a-n/a. <https://doi.org/10.1002/jbm.a.34767>
38. Kaldenhoff E, Klinge U, Klosterhalfen B, et al (2013)
Von der Prolaps- zur Problempatientin: Schenken wir der Qualität von Netzimplantaten genügend Aufmerksamkeit?
Der Gynäkologe 46:469–476. <https://doi.org/10.1007/s00129-012-3124-4>
53. Zhu L-M, Schuster P, Klinge U (2015)
An overview of crucial mesh parameters.
World Journal of Gastrointestinal Surgery
102. Klinge U, Park J-K, Klosterhalfen B (2013)
The Ideal Mesh.
Pathobiology 80:169–175. <https://doi.org/10.1159/000348446>
103. Klosterhalfen B, Klinge U (2013)
Retrieval study at 623 human mesh explants made of polypropylene - impact of mesh class and indication for mesh removal on tissue reaction.
Journal of Biomedical Materials Research Part B: Applied Biomaterials n/a-n/a. <https://doi.org/10.1002/jbmb.32958>
104. Klinge U, Junge K, Spellerberg B, et al (2002)
Do multifilament alloplastic meshes increase the infection rate? Analysis of the polymeric surface, the bacteria adherence, and the in vivo consequences in a rat model.
J Biomed Mater Res 63:765–771. <https://doi.org/10.1002/jbm.10449>
105. Klinge U, Klosterhalfen B, Birkenhauer V, et al (2002)
Impact of polymer pore size on the interface scar formation in a rat model.
J Surg Res 103:208–214. <https://doi.org/10.1006/jsre.2002.6358>

References



DynaMesh® visible


7. Hansen NL, Barabasch A, Distelmaier M, et al (2013)
First In-Human Magnetic Resonance Visualization of Surgical Mesh Implants for Inguinal Hernia Treatment.
Invest Radiol. <https://doi.org/10.1097/RLI.0b013e31829806ce>
29. Kuehnert N, Kraemer NA, Otto J, et al (2011)
In vivo MRI visualization of mesh shrinkage using surgical implants loaded with superparamagnetic iron oxides.
Surgical Endoscopy 26:1468-1475. <https://doi.org/10.1007/s00464-011-2057-7>
51. Köhler G, Pallwein-Prettner L, Lechner M, et al (2015)
First human magnetic resonance visualisation of prosthetics for laparoscopic large hiatal hernia repair.
Hernia 19:975-982. <https://doi.org/10.1007/s10029-015-1398-x>
54. Muysoms F, Beckers R, Kyle-Leinhase I (2018)
Prospective cohort study on mesh shrinkage measured with MRI after laparoscopic ventral hernia repair with an intraperitoneal iron oxide-loaded PVDF mesh.
Surgical Endoscopy 32:2822-2830. <https://doi.org/10.1007/s00464-017-5987-x>
56. Köhler G, Pallwein-Prettner L, Koch OO, et al (2015)
Magnetic Resonance-Visible Meshes for Laparoscopic Ventral Hernia Repair.
JSLS : Journal of the Society of Laparoendoscopic Surgeons 19:e2014.00175. <https://doi.org/10.4293/JSLS.2014.00175>
62. Köhler G, Wundsam H, Pallwein-Prettner L, et al (2015)
Magnetic resonance visible 3-D funnel meshes for laparoscopic parastomal hernia prevention and treatment.
European Surgery 47:127-132. <https://doi.org/10.1007/s10353-015-0319-7>
69. Kuehnert N, Otto J, Conze J, et al (2014)
Time-Dependent Changes of Magnetic Resonance Imaging-Visible Mesh Implants in Patients
70. Hansen NL, Cirlitsis A, Otto J, et al (2015)
Utility of Magnetic Resonance Imaging to Monitor Surgical Meshes: Correlating Imaging and Clinical Outcome of Patients Undergoing Inguinal Hernia Repair.
Invest Radiol. <https://doi.org/10.1097/RLI.0000000000000148>
71. Weyhe D, Klinge U, Uslar VN, et al (2019)
Follow Up Data of MRI-Visible Synthetic Meshes for Reinforcement in Large Hiatal Hernia in Comparison to None-Mesh Repair – A Prospective Cohort Study.
Front Surg 6:. <https://doi.org/10.3389/fsurg.2019.00017>
76. Lechner M, Meissnitzer M, Borhanian K, et al (2019)
Surgical and radiological behavior of MRI-depictable mesh implants after TAPP repair: the IRONMAN study.
Hernia. <https://doi.org/10.1007/s10029-019-02019-2>
90. Özveri E, Şanlı DET, Yıldırım D, et al (2020)
Magnetic resonance visualization of iron-loaded meshes in patients with pain after inguinal hernia repair.
Hernia. <https://doi.org/10.1007/s10029-020-02168-9>

References

Implant Fixation

79. Villalobos RN, Mias MC, Gas C, et al (2019)
Atraumatic laparoscopic intraperitoneal mesh fixation using a new laparoscopic device: an animal experimental study.
Hernia. <https://doi.org/10.1007/s10029-019-02008-5>
86. Wilson P (2020)
Laparoscopic intraperitoneal onlay mesh (IPOM) repair using n-butyl-2-cyanoacrylate (Liquiband Fix8TM) for mesh fixation: learning experience and short-medium term results.
Hernia. <https://doi.org/10.1007/s10029-020-02144-3>
97. Carus T (2021)
Die laparoskopische IPOM-Operation bei Nabel- und Bauchwandhernien – Netzfixierung in Klebetechnik.
6


References

Hernias

Inguinal Hernia

16. Junge K, Binnebösel M, Kauffmann C, et al (2010)
Damage to the spermatic cord by the Lichtenstein and TAPP procedures in a pig model.
Surgical Endoscopy 25:146–152. <https://doi.org/10.1007/s00464-010-1148-1>
67. Garcia-Pastor P, Porrero-Carro J, et al. (2018)
Prospective Multicenter Blinded Randomized Study Comparing PP and PVDF Mesh Implants in Lichtenstein Procedure with Respect to Pain and Recurrence.
JSM Surgical Procedures 1:
72. Guadalajara Jurado JF, Suárez Grau JM, Bellido Luque JA, et al (2016)
Initial experience in laparoscopic bilateral inguinal hernia repair (TEP) with new anatomical mesh with large pore and low weight (Dynamesh Endolap) in short stay (6 months follow-up).
Ambulatory Surgery 22:
90. Özveri E, Şanlı DET, Yıldırım D, et al (2020)
Magnetic resonance visualization of iron-loaded meshes in patients with pain after inguinal hernia repair.
Hernia. <https://doi.org/10.1007/s10029-020-02168-9>
94. Ramser M, Baur J, Keller N, et al (2021)
Robotische Hernienchirurgie: Teil I: Robotische Leistenhernienversorgung (r TAPP). Videobeitrag und Ergebnisse einer Kohortenstudie an 302 operierten Hernien.
Chirurg. <https://doi.org/10.1007/s00104-021-01425-6>
101. The HerniaSurge Group (2018)
International guidelines for groin hernia management.
Hernia. <https://doi.org/10.1007/s10029-017-1668-x>


References

Hernias

Abdominal Wall Hernia Repair

10. Berger D, Bientzle M (2008)
Polyvinylidene fluoride: a suitable mesh material for laparoscopic incisional and parastomal hernia repair!
Hernia 13:167-172. <https://doi.org/10.1007/s10029-008-0435-4>
14. Berger D, Bientzle M (2006)
Principles of laparoscopic repair of ventral hernias.
European Surgery 38:393-398. <https://doi.org/10.1007/s10353-006-0284-2>
54. Muysoms F, Beckers R, Kyle-Leinhase I (2018)
Prospective cohort study on mesh shrinkage measured with MRI after laparoscopic ventral hernia repair with an intraperitoneal iron oxide-loaded PVDF mesh.
Surgical Endoscopy 32:2822-2830. <https://doi.org/10.1007/s00464-017-5987-x>
56. Köhler G, Pallwein-Prettner L, Koch OO, et al (2015)
Magnetic Resonance-Visible Meshes for Laparoscopic Ventral Hernia Repair.
JSLS : Journal of the Society of Laparoendoscopic Surgeons 19:e2014.00175. <https://doi.org/10.4293/JSLS.2014.00175>
58. Verbo A, Pafundi P, Manno A, et al (2016)
Polyvinylidene Fluoride Mesh (PVDF, DynaMesh®-IPOM) in The Laparoscopic Treatment of Incisional Hernia: A Prospective Comparative Trial versus Gore® ePTFE DUALMESH® Plus.
Surgical technology international 28:147-151
87. Sánchez-Arteaga A, Tallón-Aguilar L, Tinoco-González J, et al (2020)
Use of polyvinylidene fluoride (PVDF) meshes for ventral hernia repair in emergency surgery.
Hernia. <https://doi.org/10.1007/s10029-020-02209-3>


References

Hernias

Abdominal Wall Hernia Prevention


55. Kohler A, Lavanchy JL, Lenoir U, et al (2019)
Effectiveness of Prophylactic Intraperitoneal Mesh Implantation for Prevention of Incisional Hernia in Patients Undergoing Open Abdominal Surgery: A Randomized Clinical Trial.
JAMA Surgery 154:109. <https://doi.org/10.1001/jamasurg.2018.4221>
74. Bravo-Salva A, González-Castillo AM, Vela-Polanco FF, et al (2019)
Incidence of Incisional Hernia After Emergency Subcostal Unilateral Laparotomy: Does Augmentation Prophylaxis Play a Role?
World J Surg. <https://doi.org/10.1007/s00268-019-05282-7>
89. Pereira JA, Pera M, López-Cano M, et al (2019)
Hernias at the Extraction Incision After Laparoscopic Colon and Rectal Resection: Influence of Incision Location and Use of Prophylactic Mesh.
Cirugía Española (English Edition) 97:20-26

References

Hernias

Parastomal Hernia Repair

9. Berger D, Bientzle M (2007)
Laparoscopic Repair of Parastomal Hernias: A Single Surgeon's Experience in 66 Patients.
Diseases of the Colon & Rectum 50:1668-1673. <https://doi.org/10.1007/s10350-007-9028-z>
12. Berger D (2010)
Laparoskopische Reparation der parastomalen Hernie.
Der Chirurg 81:988-992. <https://doi.org/10.1007/s00104-010-1933-3>
60. Fischer I, Wundsam H, Mitteregger M, Köhler G (2017)
Parastomal Hernia Repair with a 3D Funnel Intraperitoneal Mesh Device and Same-Sided Stoma Relocation: Results of 56 Cases.
World Journal of Surgery 41:3212-3217. <https://doi.org/10.1007/s00268-017-4130-4>
62. Köhler G, Wundsam H, Pallwein-Prettner L, et al (2015)
Magnetic resonance visible 3-D funnel meshes for laparoscopic parastomal hernia prevention and treatment.
European Surgery 47:127-132. <https://doi.org/10.1007/s10353-015-0319-7>
63. Köhler G, Emmanuel K (2017)
Laparoscopic stoma relocation for parastomal hernia treatment by using a magnetic resonance visible three-dimensional implant.
ANZ Journal of Surgery 87:411-412. <https://doi.org/10.1111/ans.12899>
64. Köhler G, Fischer I, Wundsam H (2018)
A Novel Technique for Parastomal Hernia Repair Combining a Laparoscopic and Ostomy-Opening Approach.
Journal of Laparoendoscopic & Advanced Surgical Techniques 28:209-214.
<https://doi.org/10.1089/jlap.2017.0313>
65. Köhler G, Mayer F, Wundsam H, et al (2015)
Changes in the Surgical Management of Parastomal Hernias Over 15 Years: Results of 135 Cases.
World Journal of Surgery 39:2795-2804. <https://doi.org/10.1007/s00268-015-3187-1>
66. Zhang H, Xie J, Miao J, Wu H (2016)
Hybrid Approaches for Complex Parastomal Hernia Repair.
Journal of the College of Physicians and Surgeons Pakistan 26:72-73
75. Köhler G (2019)
Prinzipien und Parallelen der Prävention und Reparation parastomaler Hernien mit Netzen.
Chirurg. <https://doi.org/10.1007/s00104-019-01047-z>
77. Szczepkowski M, Skoneczny P, Przywózka A, et al (2015)
Leading article: Methods paper New minimally invasive technique of parastomal hernia repair - methods and review.
witm 1:1-7. <https://doi.org/10.5114/witm.2015.50052>
78. Tully KH, Roghmann F, Pastor J, et al (2019)
Parastomal Hernia Repair With 3-D Mesh Implants After Radical Cystectomy and Ileal Conduit Urinary Diversion - A Single-center Experience Using a Purpose Made Alloplastic Mesh Implant.
Urology 131:245-249. <https://doi.org/10.1016/j.urology.2019.05.006>




Hernias

Parastomal Hernia Repair

92. Bustos-Jiménez M, Martín-Cartes JA (2020)
Surgical Treatment of Parastomal Hernias by Using A 3D Mesh.
05:6
96. Cartes JAM, Bustos-Jiménez M, Tamayo-López MJ (2018)
Parastomal Hernia: A More and More Frequent Surgical Challenge.
General Surgery 3:5


References

Hernias

Parastomal Hernia Prevention

15. Berger D (2007)
Prevention of parastomal hernias by prophylactic use of a specially designed intraperitoneal onlay mesh (Dynamesh IPST®).
Hernia 12:243–246. <https://doi.org/10.1007/s10029-007-0318-0>
59. Conde-Muñoz R, Díez J-L, Martínez A, et al (2017)
Preventing parastomal hernias with systematic intraperitoneal specifically designed mesh.
BMC Surgery 17:. <https://doi.org/10.1186/s12893-017-0237-7>
61. Köhler G, Hofmann A, Lechner M, et al (2016)
Prevention of parastomal hernias with 3D funnel meshes in intraperitoneal onlay position by placement during initial stoma formation.
Hernia 20:151–159. <https://doi.org/10.1007/s10029-015-1380-7>
62. Köhler G, Wundsam H, Pallwein-Prettner L, et al (2015)
Magnetic resonance visible 3-D funnel meshes for laparoscopic parastomal hernia prevention and treatment.
European Surgery 47:127–132. <https://doi.org/10.1007/s10353-015-0319-7>
75. Köhler G (2019)
Prinzipien und Parallelen der Prävention und Reparation parastomaler Hernien mit Netzen.
Chirurg. <https://doi.org/10.1007/s00104-019-01047-z>
80. López-Borao J, Madrazo-González Z, Kreisler E, Biondo S (2019)
Prevention of parastomal hernia after abdominoperineal excision with a prophylactic three-dimensional funnel mesh.
Colorectal Dis 21:1326–1334. <https://doi.org/10.1111/codi.14738>
88. Mäkäräinen-Uhlbäck EJ, Klintrup KHB, Vierimaa MT, et al (2020)
Prospective, Randomized Study on the Use of Prosthetic Mesh to Prevent a Parastomal Hernia in a Permanent Colostomy: Results of a Long-term Follow-up.
Diseases of the Colon & Rectum 63:678–684. <https://doi.org/10.1097/DCR.0000000000001599>
95. Ammann Y, Widmann B, Sparn M, et al (2021)
Prophylactic Funnel Mesh to Prevent Parastomal Hernia in Permanent End Colostomy: A Retrospective Cohort Study.
Colorectal Dis. <https://doi.org/10.1111/codi.15817>

References


Hernias

Hiatal Hernia Repair

51. Köhler G, Pallwein-Prettner L, Lechner M, et al (2015)
First human magnetic resonance visualisation of prosthetics for laparoscopic large hiatal hernia repair.
Hernia 19:975–982. <https://doi.org/10.1007/s10029-015-1398-x>


71. Weyhe D, Klinge U, Uslar VN, et al (2019)
Follow Up Data of MRI-Visible Synthetic Meshes for Reinforcement in Large Hiatal Hernia in Comparison to None-Mesh Repair – A Prospective Cohort Study.
Front Surg 6:.. <https://doi.org/10.3389/fsurg.2019.00017>

References

Female Pelvic Organ Prolapse

DynaMesh®-PR / -PRR

57. Jan H, Ghai V, Thakar R (2018)
Simplified Laparoscopic Sacrohysteropexy.
Journal of Minimally Invasive Gynecology 25:1134. <https://doi.org/10.1016/j.jmig.2018.01.014>

Female Pelvic Organ Prolapse

DynaMesh®-PRS

42. Balsamo R, Illiano E, Zucchi A, et al (2018)

**Sacrocolpopexy with polyvinylidene fluoride mesh for pelvic organ prolapse:
Mid term comparative outcomes with polypropylene mesh.**

European Journal of Obstetrics & Gynecology and Reproductive Biology 220:74–78. <https://doi.org/10.1016/j.ejogrb.2017.11.018>

Female Pelvic Organ Prolapse

DynaMesh®-PRP

22. Noé KG, Spüntrup C, Anapolski M (2013)
Laparoscopic pectopexy: a randomised comparative clinical trial of standard laparoscopic sacral colpo-cervicopexy to the new laparoscopic pectopexy. Short-term postoperative results.
Arch Gynecol Obstet 287:275–280. <https://doi.org/10.1007/s00404-012-2536-7>
32. Noé K-G, Schiermeier S, Alkatout I, Anapolski M (2015)
Laparoscopic Pectopexy: A Prospective, Randomized, Comparative Clinical Trial of Standard Laparoscopic Sacral Colpcocervicopexy with the New Laparoscopic Pectopexy—Postoperative Results and Intermediate-Term Follow-Up in a Pilot Study.
Journal of Endourology 29:210–215. <https://doi.org/10.1089/end.2014.0413>
45. Kale A, Biler A, Terzi H, et al (2017)
Laparoscopic pectopexy: initial experience of single center with a new technique for apical prolapse surgery.
International braz j urol 43:903–909. <https://doi.org/10.1590/s1677-5538.ibju.2017.0070>
84. Noé GK, Schiermeier S, Papathemelis T, et al (2020)
Prospective international multicenter pectopexy trial: Interim results and findings post surgery.
European Journal of Obstetrics & Gynecology and Reproductive Biology 244:81–86. <https://doi.org/10.1016/j.ejogrb.2019.10.022>
98. Noé GK, Schiermeier S, Papathemelis T, et al (2021)
Prospective International Multicenter Pelvic Floor Study: Short-Term Follow-Up and Clinical Findings for Combined Pectopexy and Native Tissue Repair.
JCM 10:217. <https://doi.org/10.3390/jcm10020217>

Female Pelvic Organ Prolapse

DynaMesh®-CESA / -VASA


33. Jaeger et al. (2016)
Does the Patients Age have an Influence on the Outcome of Cesa (Cervico-Sacropexy) and Vasa (Vagino-Sacropexy) for the Treatment of Urinary Incontinence in Women?
/ Open Access | OMICS International
34. Rajshekhar S, Mukhopadhyay S, Morris E (2016)
Early safety and efficacy outcomes of a novel technique of sacrocolpopexy for the treatment of apical prolapse.
International Journal of Gynecology and Obstetrics 0: <https://doi.org/10.1016/j.ijgo.2016.05.007>
35. Ludwig S, Stumm M, Mallmann P, Jager W (2016)
Surgical replacement of the uterosacral-and pubourethral-ligaments as treatment for urgency urinary incontinence.
Austin J Womens Health 3:1019
36. Joukhadar R, Meyberg-Solomayer G, Hamza A, et al (2015)
A Novel Operative Procedure for Pelvic Organ Prolapse Utilizing a MRI-Visible Mesh Implant: Safety and Outcome of Modified Laparoscopic Bilateral Sacropexy.
BioMed Research International 2015:1-9. <https://doi.org/10.1155/2015/860784>
73. Cassis C, Mukhopadhyay S, Morris E (2019)
Standardizing abdominal sacrocolpopexy for the treatment of apical prolapse: One year on.
Int J Gynecol Obstet ijgo.12935. <https://doi.org/10.1002/ijgo.12935>
85. Rexhepi S, Rexhepi E, Stumm M, et al (2018)
Laparoscopic Bilateral Cervicosacropexy and Vaginosacropexy: New Surgical Treatment Option in Women with Pelvic Organ Prolapse and Urinary Incontinence.
Journal of Endourology 32:1058-1064. <https://doi.org/10.1089/end.2018.0474>

References

Female Urinary Incontinence

21. Klinge U, Binneboesel M, Kuschel S, Schuessler B (2007)
Demands and properties of alloplastic implants for the treatment of stress urinary incontinence.
Expert Rev Med Devices 4:349–359. <https://doi.org/10.1586/17434440.4.3.349>
31. Naumann G, Albrich S, Skala C, et al (2012)
Single-Incision Slings (SIS) - A New Option for the Surgical Treatment of Female Stress Urinary Incontinence.
Geburtshilfe und Frauenheilkunde 72:125–131. <https://doi.org/10.1055/s-0031-1298275>
39. Ludwig S, Stumm M, Mallmann P, Jager W (2016)
TOT 8/4: A Way to Standardize the Surgical Procedure of a Transobturator Tape.
BioMed Research International 2016:1–4. <https://doi.org/10.1155/2016/4941304>
40. Najjari L, Hennemann J, Kirschner-Hermanns R, et al (2014)
Visualization of Polypropylene and Polyvinylidene Fluoride Slings in Perineal Ultrasound and Correlation with Clinical Outcome.
BioMed Research International. <https://doi.org/10.1155/2014/181035>
41. Sabadell J, Larrain F, Gracia-Perez-Bonfils A, et al (2016)
Comparative study of polyvinylidene fluoride and polypropylene suburethral slings in the treatment of female stress urinary incontinence: PVDF/polypropylene in suburethral slings.
Journal of Obstetrics and Gynaecology Research 42:291–296. <https://doi.org/10.1111/jog.12899>
44. Ludwig S, Stumm M (2016)
Surgical Treatment of Urgency Urinary Incontinence, OAB (Wet), Mixed Urinary Incontinence, and Total Incontinence by Cervicosacropexy or Vaginosacropexy.
Gynecology & Obstetrics 6:.. <https://doi.org/10.4172/2161-0932.1000404>
49. Najjari L, Gräf CM, Kupec T, et al (2016)
Tomographic Ultrasound Imaging to Control the Placement of Tension-Free Transobturator Tape in Female Urinary Stress Incontinence.
BioMed Research International 2016:1–6. <https://doi.org/10.1155/2016/6495858>
83. Ludwig S, Becker I, Mallmann P, Jäger W (2019)
Comparison of Solifenacin and Bilateral Apical Fixation in the Treatment of Mixed and Urgency Urinary Incontinence in Women: URGE 1 Study, A Randomized Clinical Trial.
In Vivo 33:1949–1957. <https://doi.org/10.21873/invivo.11690>
99. Sabadell J, Pereda-Núñez A, Ojeda-de-los-Santos F, et al (2021)
Polypropylene and polyvinylidene fluoride transobturator slings for the treatment of female stress urinary incontinence: 1-Year outcomes from a multicentre randomized trial.
Neurourology and Urodynamics 40:475–482. <https://doi.org/10.1002/nau.24586>

References

Male Urinary Incontinence

81. Costa Cruz DSL da, D'Ancona CAL, Silva Filho WP da, et al (2020)
Parameters of 2-Dimensional Perineal Ultrasonography Before and After Male Sling Procedure for Urinary Incontinence After Radical Prostatectomy.
Urology 136:257-262. <https://doi.org/10.1016/j.urology.2019.10.004>

Internal test-report references

Internal test-reports

TR1. CF_F02 "Biocompatibility"

TR10. I1_F03-01_SIS1_X Stability and Elongation (*bench test*)TR11. I1_F03-02_SIS1_X Porosity and Formstability (*bench test*)TR12. I1_F03-03_SIS1_X Formstability (*bench test*)